



Author index

Volume 38 (2005)

- Alper, S., see Itrli, G. 38, 225
 Atsuko, M., see Masako, K. 38, 207
- Baba, H., M. Yoshida, T. Yokota, H. Uchiwa, S. Watanabe, Human epidermal basal cell responses to ultraviolet-B differ according to their location in the undulating epidermis 38, 41
 Bacon, P., B. Bodner, B.J. Nickoloff, Senescent human keratinocytes suppress colony formation of HeLa cells 38, 64
 Bavinck, J.N.B., see de Graaf, Y.G.L. 38, 228
 Benkő, R., see Racz, E. 38, 231
 Bodner, B., see Bacon, P. 38, 64
- Cai, X.N., see Li, Y. 38, 107
 Caproni, M., see Giomi, B. 38, 57
 Chan, I., see Kowalewski, C. 38, 215
 Chang, S.-Y., see Rho, S.-S. 38, 89
 Cho, B.K., see Park, H.J. 38, 9
 Cho, D.H., see Park, H.J. 38, 9
 Csikós, M., see Racz, E. 38, 231
- de Graaf, Y.G.L., B.J. Vermeer, J.N.B. Bavinck, R. Willemze, F.R. de Gruijl, D. Schiefer, A. Redeker, S.H. Van der Burg, p53-Specific serum antibodies are not associated with a history of skin carcinoma in renal transplant recipients and immunocompetent individuals 38, 228
 de Gruijl, F.R., see de Graaf, Y.G.L. 38, 228
- Fabbri, P., see Giomi, B. 38, 57
 Fan, J.-Y., see Shen, Z. 38, 25
 Fujimoto, W., see Uchida, T. 38, 60
- Giomi, B., M. Caproni, P. Fabbri, IL-4 and cellular adhesion molecule (CAM) pathway are involved in cicatricial pemphigoid scarring process 38, 57
 Górski, M., see Kowalewski, C. 38, 215
 Goto, I., see Kanbe, T. 38, 99
- Hideyuki, I., see Masako, K. 38, 197, 207
 Hino, R., T. Shimauchi, Y. Tokura, Treatment with IFN- γ increases serum levels of Th1 chemokines and decreases those of Th2 chemokines in patients with mycosis fungoides 38, 189
 Hwang, S.-L., see Rho, S.-S. 38, 89
- Iino, M., see Soma, T. 38, 110
 Inaoki, M., see Uchida, T. 38, 60
 Ishizaki, H., see Kanbe, T. 38, 99
- Itrli, G., M. Pehlivan, S. Alper, S.E. Yüksel, H. Onay, F. Özkinay, S. Pehlivan, Exon-3 polymorphism of CTLA-4 gene in Turkish patients with vitiligo 38, 225
- Jabłońska, S., see Kowalewski, C. 38, 215
- Kanbe, T., L. Natsume, I. Goto, M. Kawasaki, T. Mochizuki, H. Ishizaki, A. Kikuchi, Rapid and specific identification of *Sporothrix schenckii* by PCR targeting the DNA topoisomerase II gene 38, 99
 Kanda, N., S. Watanabe, Regulatory roles of sex hormones in cutaneous biology and immunology 38, 1
 Kárpáti, S., see Racz, E. 38, 231
 Kato, T., see Matsuoka, A. 38, 17
 Kawasaki, M., see Kanbe, T. 38, 99
 Kayoko, M., see Masako, K. 38, 207
 Khera, P., see Koo, J. 38, 75
 Kikuchi, A., see Kanbe, T. 38, 99
 Kikuchi, M., see Kuwahara, K. 38, 47
 Kim, C.D., see Rho, S.-S. 38, 89
 Kim, C.W., S. Suh, S.H. Sung, I.K. Lee, K.S. Lee, Corrigendum to “A transcriptional factor decoy against AP-1 suppresses TGF- β 1-induced type I collagen gene expression in cultured keloid fibroblasts”. [J. Dermatol. Sci. 37 (2005) 49-51] 38, 71
 Kim, H.J., see Park, H.J. 38, 9
 Kim, H.S., see Park, H.J. 38, 9
 Kishimoto, J. see Toma, T. 38, 110
 Kitagaki, H., see Kuwahara, K. 38, 47
 Kitazawa, T., see Kuwahara, K. 38, 47
 Kobayashi, M., Y. Tokura, Preferential downmodulation of certain chemokines by fexofenadine in human keratinocytes 38, 67
 Koo, J., P. Khera, Update on the mechanisms and efficacy of biological therapies for psoriasis 38, 75
 Kornseé, Z., see Racz, E. 38, 231
 Kowalewski, C., A. Kozłowska, I. Chan, M. Górski, K. Woźniak, S. Jabłońska, J.A. McGrath, Three-dimensional imaging reveals major changes in skin microvasculature in lipoid proctitis and lichen sclerosis 38, 215
 Kozłowska, A., see Kowalewski, C. 38, 215
 Kuwahara, K., T. Kitazawa, H. Kitagaki, T. Tsukamoto, M. Kikuchi, Nadifloxacin, an antiacne quinolone antimicrobial, inhibits the production of proinflammatory cytokines by human peripheral blood mononuclear cells and normal human keratinocytes 38, 47
- Lee, A.W., see Park, H.J. 38, 9
 Lee, I.-H., see Rho, S.-S. 38, 89

- Lee, I.K., see Kim, C.W. 38, 71
 Lee, J.Y., see Park, H.J. 38, 9
 Lee, M.-H., see Rho, S.-S. 38, 89
 Li, G.Q., see Li, Y. 38, 107
 Li, W., see Shen, Z. 38, 25
 Li, Y., C.M. Lin, X.N. Cai, G.Q. Li, Reconstruction of human hair dermal papilla with microencapsulation in vitro 38, 107
 Lin, C.M., see Li, Y. 38, 107
 Liu, Y.-F., see Shen, Z. 38, 25
 Luger, T., see Ständer, S. 38, 177
- Makino, E., see Uchida, T. 38, 60
 Makoto, K., see Masako, K. 38, 207
 Masako, K., I. Hideyuki, O. Shigeyuki, I. Zenro, A novel method to control the balance of skin microflora. Part 1. Attack on biofilm of *Staphylococcus aureus* without antibiotics 38, 205
 Masako, K., K. Yusuke, I. Hideyuki, M. Atsuko, M. Yoshiki, M. Kayoko, K. Makoto, A novel method to control the balance of skin microflora. Part 2. A study to assess the effect of a cream containing farnesol and xylitol on atopic dry skin 38, 207
 Matsuoka, A., T. Kato, Y. Soma, H. Takahama, M. Nakamura, H. Matsuoka, M. Mizoguchi, Analysis of T cell receptor (TCR) BV-gene clonotypes in NC/Nga mice developing dermatitis resembling human atopic dermatitis 38, 17
 Matsuoka, H., see Matsuoka, A. 38, 17
 McGrath, J.A., see Kowalewski, C. 38, 215
 McLean, W.H.I., see White, S.J. 38, 169
 Metze, D., see Ständer, S. 38, 177
 Mizoguchi, M., see Matsuoka, A. 38, 17
 Mochizuki, T., see Kanbe, T. 38, 99
- Nakamura, M., see Matsuoka, A. 38, 17
 Natsume, L., see Kanbe, T. 38, 99
 Nickoloff, B.J., see Bacon, P. 38, 64
- Onay, H., see Itırlı, G. 38, 225
 Özkınay, F., see Itırlı, G. 38, 225
- Park, H.J., H.S. Kim, H.J. Kim, J.Y. Lee, B.K. Cho, A.W. Lee, D.Y. Yoon, D.H. Cho, Immunohistochemical characterization of cutaneous drug eruptions by STI571 38, 9
 Park, S.-J., see Rho, S.-S. 38, 89
 Pehlivan, M., see Itırlı, G. 38, 225
 Pehlivan, S., see Itırlı, G. 38, 225
- Racz, E., M. Csikós, R. Benkó, Z. Kornseé, S. Kárpáti, Three novel mutations in the *ATP2A2* gene in Hungarian families with Darier's disease, including a novel splice site generating intronic nucleotide change 38, 231
 Rang, M.-J., see Rho, S.-S. 38, 89
 Redeker, A., see de Graaf, Y.G.L. 38, 228
- Rho, S.-S., S.-J. Park, S.-L. Hwang, M.-H. Lee, C.D. Kim, I.-H. Lee, S.-Y. Chang, M.-J. Rang, The hair growth promoting effect of *Asiasari radix* extract and its molecular regulation 38, 89
 Rukwied, R., see Ständer, S. 38, 177
- Schiefer, D., see de Graaf, Y.G.L. 38, 228
 Schmelz, M., see Ständer, S. 38, 177
 Shen, Z., G. Wang, J.-Y. Fan, W. Li, Y.-F. Liu, HLA DR B1*04, *07-restricted epitopes on Keratin 17 for autoreactive T cells in psoriasis 38, 25
 Shigeyuki, O., see Masako, K. 38, 197
 Shimauchi, T., see Hino, R. 38, 189
 Soma, T., M. Iino, M. Tajima, J. Kishimoto, Expression of novel keratin associated protein 5 genes in the cuticle layer of human hair follicles 38, 110
 Soma, Y., see Matsuoka, A. 38, 177
 Ständer, S., M. Schmelz, D. Metze, T. Luger, R. Rukwied, Distribution of cannabinoid receptor 1 (CB1) and 2 (CB2) on sensory nerve fibers and adnexal structures in human skin 38, 177
 Suh, S., see Kim, C.W. 38, 71
 Sung, S.H., see Kim, C.W. 38, 71
- Tajima, M., see Soma, T. 38, 110
 Takahama, H., see Matsuoka, A. 38, 17
 Tokura, Y., see Hino, R. 38, 189
 Tokura, Y., see Kobayashi, M. 38, 67
 Tsukamoto, T., see Kuwahara, K. 38, 47
- Uchida, T., M. Inaoki, E. Makino, W. Fujimoto, Identification of a recurrent mutation in keratin 17 in a Japanese family with pachyonychia congenita type 2 38, 60
 Uchiwa, H., see Baba, H. 38, 41
- Van der Burg, S.H., see de Graaf, Y.G.L. 38, 228
 Vermeer, B.J., see Graaf, Y.G.L.d. 38, 228
- Wang, G., see Shen, Z. 38, 25
 Watanabe, S., see Baba, H. 38, 41
 Watanabe, S., see Kanda, N. 38, 1
 White, S.J., W.H.I. McLean, Kindler surprise: mutations in a novel actin-associated protein cause Kindler syndrome 38, 169
 Willemze, R., see de Graaf, Y.G.L. 38, 228
 Woźniak, K., see Kowalewski, C. 38, 215
- Yokota, T., see Baba, H. 38, 41
 Yoon, D.Y., see Park, H.J. 38, 9
 Yoshida, M., see Baba, H. 38, 41
 Yoshiki, M., see Masako, K. 38, 207
 Yüksel, S.E., see Itırlı, G. 38, 225
 Yusuke, K., see Masako, K. 38, 207
- Zenro, I., see Masako, K. 38, 197